

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57020 A1(51) International Patent Classification⁷: C07D 401/12,
A61K 31/4184, C07D 401/14, 475/04, 403/14, 235/16,
209/10, A61P 7/02, A61K 31/42(74) Agent: LEE, Christine, S.; Morgan, Lewis & Bockius
LLP, 1800 M. Street, N.W., Washington, DC 20036-5869
(US).

(21) International Application Number: PCT/US01/03225

(22) International Filing Date: 1 February 2001 (01.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/179,390 1 February 2000 (01.02.2000) US
60/191,735 24 March 2000 (24.03.2000) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): COR
THERAPEUTICS, INC. [US/US]; 256 E. Grand Avenue,
South Francisco, CA 94080 (US).**Published:**


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ZHU, Bing-Yan
[CA/US]; 3325 Adelaide Way, Belmont, CA 94002 (US).
SCARBOROUGH, Robert [US/US]; 22 Greenbrier
Court, Half Moon Bay, CA 94019 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INDOLE AND BENZIMIDAZOLE INHIBITORS OF FACTOR Xa



(57) Abstract: Novel compounds of formula (I) including its pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives having activity against mammalian factor Xa is described. Compositions containing such compounds are also described. The compounds and compositions are useful *in vitro* or *in vivo* for preventing or treating conditions in mammals characterized by undesired thrombosis.

INDOLE AND BENZIMIDAZOLE INHIBITORS OF FACTOR Xa

Field of the Invention

The invention relates to novel indole-containing and benzimidazole-containing compounds including their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives, and pharmaceutically acceptable compositions thereof which are potent and highly selective inhibitors of isolated factor Xa or when assembled in the prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation (e.g. thrombin, fVIIa, fIXa) or the fibrinolytic cascades (e.g. plasminogen activators, plasmin). In another aspect, the present invention relates to novel indole-containing and benzimidazole-containing compounds including their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives, and pharmaceutically acceptable compositions thereof which are useful as potent and specific inhibitors of blood coagulation in mammals. In yet another aspect, the invention relates to methods for using these inhibitors as diagnostic or therapeutic agents for disease states in mammals characterized by undesired thrombosis or coagulation disorders.

Background of the Invention

Hemostasis, the control of bleeding, occurs by surgical means, or by the physiological properties of vasoconstriction and coagulation. The invention is particularly concerned with blood coagulation and ways in which it assists in maintaining the integrity of mammalian circulation after injury, inflammation, disease, congenital defect, dysfunction or other disruption. Under normal hemostatic circumstances, the body maintains an acute balance of clot formation and clot removal (fibrinolysis). The blood coagulation cascade involves the conversion of a variety of inactive enzymes (zymogens) into active enzymes which ultimately convert the soluble plasma protein fibrinogen into an insoluble matrix of highly cross-linked fibrin. Davie, E.J. *et al.*, "The Coagulation Cascade: Initiation, Maintenance and Regulation", *Biochemistry*, 30, 10363-10370 (1991). These plasma glycoprotein zymogens include Factor XII, Factor XI, Factor IX, Factor X, Factor VII, and prothrombin. Blood coagulation follows either the intrinsic pathway, where all of the

protein components are present in blood, or the extrinsic pathway, where the cell-membrane protein tissue factor plays a critical role. Clot formation occurs when fibrinogen is cleaved by thrombin to form fibrin. Blood clots are composed of activated platelets and fibrin.

5 Blood platelets which adhere to damaged blood vessels are activated and incorporated into the clot and thus play a major role in the initial formation and stabilization of hemostatic "plugs". In certain diseases of the cardiovascular system, deviations from normal hemostasis push the balance of clot formation and clot dissolution towards life-threatening thrombus formation when thrombi occlude blood
10 flow in coronary vessels (myocardial infarctions) or limb and pulmonary veins (venous thrombosis). Although platelets and blood coagulation are both involved in thrombus formation, certain components of the coagulation cascade are primarily responsible for the amplification or acceleration of the processes involved in platelet aggregation and fibrin deposition.

15 Thrombin is a key enzyme in the coagulation cascade as well as in hemostasis. Thrombin plays a central role in thrombosis through its ability to catalyze the conversion of fibrinogen into fibrin and through its potent platelet activation activity. Under normal circumstances, thrombin can also play an anticoagulant role in hemostasis through its ability to convert protein C into
20 activated protein C (aPC) in a thrombomodulin-dependent manner. However, in atherosclerotic arteries these thrombin activities can initiate the formation of a thrombus, which is a major factor in pathogenesis of vasoocclusive conditions such as myocardial infarction, unstable angina, nonhemorrhagic stroke and reocclusion of coronary arteries after angioplasty or thrombolytic
25 therapy. Thrombin is also a potent inducer of smooth muscle cell proliferation and may therefore be involved in a variety of proliferative responses such as restenosis after angioplasty and graft induced atherosclerosis. In addition, thrombin is chemotactic for leukocytes and may therefore play a role in inflammation. (Hoover, R.J., et al. Cell , 14, 423 (1978); Etingin, O.R., et al.,
30 Cell, 61, 657 (1990). These observations indicate that inhibition of thrombin formation or inhibition of thrombin itself may be effective in preventing or treating thrombosis, limiting restenosis and controlling inflammation.

Direct or indirect inhibition of thrombin activity has been the focus of a variety of recent anticoagulant strategies as reviewed by Claeson, G., "Synthetic Peptides and Peptidomimetics as Substrates and Inhibitors of Thrombin and Other Proteases in the Blood Coagulation System", *Blood Coag. Fibrinol.* 5, 411-436 (1994). Several classes of anticoagulants currently used in the clinic directly or indirectly affect thrombin (i.e. heparins, low-molecular weight heparins, heparin-like compounds and coumarins).

The formation of thrombin is the result of the proteolytic cleavage of its precursor prothrombin at the Arg-Thr linkage at positions 271-272 and the Arg-Ile linkage at positions 320-321. This activation is catalyzed by the prothrombinase complex, which is assembled on the membrane surfaces of platelets, monocytes, and endothelial cells. The complex consists of Factor Xa (a serine protease), Factor Va (a cofactor), calcium ions and the acidic phospholipid surface. Factor Xa is the activated form of its precursor, Factor X, which is secreted by the liver as a 58 kd precursor and is converted to the active form, Factor Xa, in both the extrinsic and intrinsic blood coagulation pathways. Factor X is a member of the calcium ion binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood coagulation glycoprotein family, which also includes Factors VII and IX, prothrombin, protein C and protein S (Furie, B., et al., *Cell*, 53, 505 (1988)). The activity of Factor Xa in effecting the conversion of prothrombin to thrombin is dependent on its inclusion in the prothrombinase complex.

The prothrombinase complex converts the zymogen prothrombin into the active procoagulant thrombin. It is therefore understood that Factor Xa catalyzes the next-to-last step in the blood coagulation cascade, namely the formation of the serine protease thrombin. In turn, thrombin then acts to cleave soluble fibrinogen in the plasma to form insoluble fibrin.

The location of the prothrombinase complex at the convergence of the intrinsic and extrinsic coagulation pathways, and the resulting significant amplification of thrombin generation (several hundred-thousand fold faster in effecting the conversion of prothrombin to thrombin than Factor Xa in soluble form) mediated by the complex at a limited number of targeted catalytic units present at vascular lesion sites, suggests that inhibition of thrombin generation is a desirable method to block uncontrolled

procoagulant activity. It has been suggested that compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic agents, or for therapeutic administration in certain thrombotic disorders, see *e.g.*, WO 94/13693. Unlike thrombin, which acts on a variety of protein substrates as well as at a specific receptor, factor Xa appears to
5 have a single physiologic substrate, namely prothrombin.

Plasma contains an endogenous inhibitor of both the factor VIIa-tissue factor (TF) complex and factor Xa called tissue factor pathway inhibitor (TFPI). TFPI is a Kunitz-type protease inhibitor with three tandem Kunitz domains. TFPI inhibits the TF/fVIIa complex in a two-step mechanism which includes the initial interaction of
10 the second Kunitz domain of TFPI with the active site of factor Xa, thereby inhibiting the proteolytic activity of factor Xa. The second step involves the inhibition of the TF/fVIIa complex by formation of a quaternary complex TF/fVIIa/TFPI/fXa as described by Girard, T.J. *et al.*, "Functional Significance of the Kunitz-type Inhibitory Domains of Lipoprotein-associated Coagulation Inhibitor", *Nature*, 338, 518-520
15 (1989).

Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. United States Patent 4,588,587 describes anticoagulant activity in the saliva of the Mexican leech, *Haementeria officinalis*. A principal component of this saliva was shown to be the
20 polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. *et al.*, "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263, 10162-10167 (1988).

Another potent and highly specific inhibitor of Factor Xa, called tick anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft
25 tick *Ornithodoros moubata*, as reported by Waxman, L., *et al.*, "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa" *Science*, 248, 593-596 (1990).

Other polypeptide type inhibitors of factor Xa have been reported including the following: Condra, C. *et al.*, "Isolation and Structural Characterization of a Potent
30 Inhibitor of Coagulation Factor Xa from the Leech *Haementeria ghilianii*", *Thromb. Haemost.*, 61, 437-441 (1989); Blankenship, D.T. *et al.*, "Amino Acid Sequence of Ghilanten: Anti-coagulant-antimetastatic Principle of the South American Leech,

- Haementeria ghilianii*", Biochem. Biophys. Res. Commun. 166, 1384-1389 (1990); Brankamp, R.G. *et al.*, "Ghilantens: Anticoagulants, Antimetastatic Proteins from the South American Leech *Haementeria ghilianii*", J. Lab. Clin. Med., 115, 89-97 (1990); Jacobs, J.W. *et al.*, "Isolation and Characterization of a Coagulation Factor Xa Inhibitor from Black Fly Salivary Glands", Thromb. Haemost., 64, 235-238 (1990); Rigbi, M. *et al.*, "Bovine Factor Xa Inhibiting Factor and Pharmaceutical Compositions Containing the Same", European Patent Application, 352,903; Cox, A.C., "Coagulation Factor X Inhibitor From the Hundred-pace Snake *Deinagkistrodon acutus*, venom", Toxicon, 31, 1445-1457 (1993); Cappello, M. *et al.*, "Ancylostoma Factor Xa Inhibitor: Partial Purification and its Identification as a Major Hookworm-derived Anticoagulant In Vitro", J. Infect. Dis., 167, 1474-1477 (1993); Seymour, J.L. *et al.*, "Ecotin is a Potent Anticoagulant and Reversible Tight-binding Inhibitor of Factor Xa", Biochemistry 33, 3949-3958 (1994).

- Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors have also been reported including: Tidwell, R.R. *et al.*, "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin Inhibitors", Thromb. Res., 19, 339-349 (1980); Turner, A.D. *et al.*, "p-Amidino Esters as Irreversible Inhibitors of Factor IXa and Xa and Thrombin", Biochemistry, 25, 4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", Haemostasis, 15, 164-168 (1985); Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", Thromb. Res., 54, 245-252 (1989); Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", Biochemistry, 27, 2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic Effects of Synthetic Thrombin and Factor Xa Inhibitors", Thromb. Haemost., 63, 220-223 (1990); Miyadera, A. *et al.*, Japanese Patent Application JP 6327488; Nagahara, T. *et al.*, "Dibasic (Amidinoaryl)propanoic Acid Derivatives as Novel Blood Coagulation Factor Xa Inhibitors", J. Med. Chem., 37, 1200-1207 (1994); Vlasuk, G.P. *et al.*, "Inhibitors of Thrombosis", WO 93/15756; and Brunck, T.K. *et al.*, "Novel Inhibitors of Factor Xa", WO 94/13693.

A number of inhibitors of trypsin-like enzymes (such as trypsin, enterokinase, thrombin, kallikrein, plasmin, urokinase, plasminogen activators and the like) have been the subject of disclosures. For example, Austen *et al.*, United States Patent 4,593,018 describes oligopeptide aldehydes which are specific inhibitors of enterokinase; Abe *et al.*, United States Patent 5,153,176 describes tripeptide aldehydes which have inhibitory activity against multiple serine proteases such as plasmin, thrombin, trypsin, kallikrein, factor Xa, urokinase, etc.; Brunck *et al.*, European Publication WO 93/14779 describes substituted tripeptide aldehydes that are specific inhibitors of trypsin; United States Patents 4,316,889, United States Patent 4,399,065, United States Patent 4,478,745 all disclose arginine aldehyde inhibitors of thrombin; Balasubramanian *et al.*, United States Patent 5,380,713 describes di and tripeptide aldehydes which are useful for anti-trypsin and anti-thrombin activity; Webb *et al.*, United States Patent 5,371,072 describes tripeptide alpha-keto-amide derivatives as inhibitors of thrombosis and thrombin; Gesellchen *et al.*, European Patent Publications 0479489 A2 and 0643073 A, describe tripeptide thrombin inhibitors; Veber *et al.*, European Publication WO 94/25051 describes 4-cyclohexylamine derivatives which selectively inhibit thrombin over other trypsin-like enzymes; Tapparelli *et al.*, J. Biol. Chem. 268, 4734-4741 (1993) describe selective peptide boronic acid derivatives as inhibitors of thrombin.

Alternatively, agents which inhibit the vitamin K-dependent carboxylase enzyme, such as coumarin, have been used to treat coagulation disorders.


There exists a need for effective therapeutic agents for the regulation of hemostasis, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as restenosis and inflammation.

Summary of the Invention

The present invention provides novel indole-containing and benzimidazole-containing compounds including their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives, which have particular biological properties and are useful as potent and specific inhibitors of blood coagulation in mammals. The invention also provides compositions containing such compounds. The compounds of

the invention may be used as diagnostic reagents or as therapeutic agents for disease states in mammals which have coagulation disorders. Thus, the invention further provides methods for preventing or treating a condition in a mammal characterized by undesired thrombosis by administration of a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. Optionally, the methods of the invention comprise administering a pharmaceutical composition of the invention in combination with an additional therapeutic agent such as an antithrombotic and/or a thrombolytic agent and/or an anticoagulant. According to the invention, such conditions include, for example, any thrombotically mediated acute coronary or cerebrovascular syndrome, any thrombotic syndrome occurring in the venous system, any coagulopathy, and any thrombotic complications associated with extracorporeal circulation or instrumentation. The invention still further provides a method for inhibiting the coagulation of biological samples (e.g. stored blood products and samples).


The invention provides a compound of general formula I:



wherein:

8

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$;
 $-C(=O)NR^3R^4$;



5

where R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and
 10 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

15

m is an integer from 0-3, preferably 0-2;

Z is a member selected from the group consisting of a direct link, C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group
 20 consisting of N, O and S;

n is an integer from 0-3, preferably 0-2;

D is a member selected from the group consisting of a direct link, $-CH_2-$, $-O-$,
 25 $-N(R^2)-$, $-C(=O)-$, $-S-$, $-SO_2-$, $-SO_2-N(R^2)-$, $-N(R^2)-SO_2-$, $-OC(=O)-$, $-C(=O)O-$,

-C(=O)-N(R²)- and -N(R²)-C(=O)-, where R² is as described above;

- R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH,
- 5 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and
- 10 C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² and R³ are as described above;

q is an integer from 0-3, preferably 0-2;

- R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl,
- 15 C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
- 20 form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

X is N or -CR¹¹; where R¹¹ is defined as above;

25


p is an integer from 0-3, preferably 0-2;

- E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)-, where R¹¹ is as defined above, phenylene, a bivalent 5 to 12 member
- 30 bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group

consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups and each R^{14} group is independently defined the same as the substituents set forth above for the R^1 group;

- 5 Q is a member selected from the group consisting of a direct link, a bivalent C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by
- 10 from 0 to 5 R^{14} groups and each R^{14} group is independently defined the same as the substituents set forth above for the R^1 group;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



15

wherein

t is an integer from 0 to 6,

u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are

independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

Detailed Description of the InventionDefinitions

In accordance with the present invention and as used herein, the following terms are defined with the following meanings, unless explicitly stated otherwise.

5 The term "alkenyl" refers to a trivalent straight chain or branched chain unsaturated aliphatic radical. The term "alkynyl" (or "alkynyl") refers to a straight or branched chain aliphatic radical that includes at least two carbons joined by a triple bond. If no number of carbons is specified alkenyl and alkynyl each refer to radicals having from 2-12 carbon atoms.

10 The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain and cyclic groups having the number of carbon atoms specified, or if no number is specified, having up to 12 carbon atoms. The term "lower alkyl" refers to a C₁-C₈ unsubstituted alkyl group unless a substituent(s) is specified. The term "cycloalkyl" as used herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3
15 to 14 carbon atoms and preferably 3 to 7 carbon atoms.

As used herein, the terms "carbocyclic ring structure" and "C₃₋₁₆ carbocyclic mono, bicyclic or tricyclic ring structure" or the like are each intended to mean stable ring structures having only carbon atoms as ring atoms wherein the ring structure is a substituted or unsubstituted member selected from the group consisting of: a stable
20 monocyclic ring which is aromatic ring ("aryl") having six ring atoms; a stable monocyclic non-aromatic ring having from 3 to 7 ring atoms in the ring; a stable bicyclic ring structure having a total of from 7 to 12 ring atoms in the two rings wherein the bicyclic ring structure is selected from the group consisting of ring structures in which both of the rings are aromatic, ring structures in which one of the
25 rings is aromatic and ring structures in which both of the rings are non-aromatic; and a stable tricyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein the tricyclic ring structure is selected from the group consisting of: ring structures in which three of the rings are aromatic, ring structures in which two of the rings are aromatic and ring structures in which three of the rings are non-aromatic. In
30 each case, the non-aromatic rings when present in the monocyclic, bicyclic or tricyclic

ring structure may independently be saturated, partially saturated or fully saturated.

Examples of such carbocyclic ring structures include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin),

- 5 2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any carbon atom which results in a stable structure. The term "substituted" as used in conjunction with carbocyclic ring structures means that hydrogen atoms attached to the ring carbon atoms of ring
- 10 structures described herein may be substituted by one or more of the substituents indicated for that structure if such substitution(s) would result in a stable compound.

- The term "aryl" which is included with the term "carbocyclic ring structure" refers to an unsubstituted or substituted aromatic ring, substituted with one, two or three substituents selected from loweralkoxy, loweralkyl, loweralkylamino, hydroxy,
- 15 halogen, cyano, hydroxyl, mercapto, nitro, thioalkoxy, carboxaldehyde, carboxyl, carboalkoxy and carboxamide, including but not limited to carbocyclic aryl, heterocyclic aryl, and biaryl groups and the like, all of which may be optionally substituted. Preferred aryl groups include phenyl, halophenyl, loweralkylphenyl, naphthyl, biphenyl, phenanthrenyl and naphthacenyl.

- 20 The term "arylalkyl" which is included with the term "carbocyclic aryl" refers to one, two, or three aryl groups having the number of carbon atoms designated, appended to an alkyl group having the number of carbon atoms designated. Suitable arylalkyl groups include, but are not limited to, benzyl, picolyl, naphthylmethyl, phenethyl, benzyldryl, trityl, and the like, all of which may be optionally substituted.

- 25 As used herein, the term "heterocyclic ring" or "heterocyclic ring system" is intended to mean a substituted or unsubstituted member selected from the group consisting of stable monocyclic ring having from 5-7 members in the ring itself and having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S; a stable bicyclic ring structure having a total of from 7 to 12 atoms in the two rings
- 30 wherein at least one of the two rings has from 1 to 4 hetero atoms selected from N, O

and S, including bicyclic ring structures wherein any of the described stable monocyclic heterocyclic rings is fused to a hexane or benzene ring; and a stable tricyclic heterocyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein at least one of the three rings has from 1 to 4 hetero atoms selected from the group consisting of N, O and S. Any nitrogen and sulfur atoms present in a heterocyclic ring of such a heterocyclic ring structure may be oxidized. Unless indicated otherwise the terms "heterocyclic ring" or "heterocyclic ring system" include aromatic rings, as well as non-aromatic rings which can be saturated, partially saturated or fully saturated non-aromatic rings. Also, unless indicated otherwise the term "heterocyclic ring system" includes ring structures wherein all of the rings contain at least one hetero atom as well as structures having less than all of the rings in the ring structure containing at least one hetero atom, for example bicyclic ring structures wherein one ring is a benzene ring and one of the rings has one or more hetero atoms are included within the term "heterocyclic ring systems" as well as bicyclic ring structures wherein each of the two rings has at least one hetero atom. Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any hetero atom or carbon atom which results in a stable structure. Further, the term "substituted" means that one or more of the hydrogen atoms on the ring carbon atom(s) or nitrogen atom(s) of the each of the rings in the ring structures described herein may be replaced by one or more of the indicated substituents if such replacement(s) would result in a stable compound. Nitrogen atoms in a ring structure may be quaternized, but such compounds are specifically indicated or are included within the term "a pharmaceutically acceptable salt" for a particular compound. When the total number of O and S atoms in a single heterocyclic ring is greater than 1, it is preferred that such atoms not be adjacent to one another. Preferably, there are no more than 1 O or S ring atoms in the same ring of a given heterocyclic ring structure.

Examples of monocyclic and bicyclic heterocyclic ring systems, in alphabetical order, are acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl,

2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, 1H-indazolyl, indoliny, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindoliny, isoindolyl, isoquinoliny (benzimidazolyl), isothiazolyl, isoxazolyl, morpholiny, naphthyridiny, 5 octahydroisoquinoliny, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidiny, oxazolyl, oxazolidiny, pyrimidiny, phenanthridiny, phenanthroliny, phenaziny, phenothiaziny, phenoxathiiny, phenoxaziny, phthalaziny, piperaziny, piperidiny, pteridiny, puriny, pyranyl, pyraziny, pyroazolidiny, pyrazoliny, pyrazolyl, pyridaziny, 10 pryidooxazole, pyridoimidazole, pyridothiazole, pyridiny, pyridyl, pyrimidiny, pyrrolidiny, pyrroliny, 2H-pyrroly, pyrroly, quinazoliny, quinoliny, 4H-quinoliziny, quinoxaliny, quinuclidiny, tetrahydrofuranyl, tetrahydroisoquinoliny, tetrahydroquinoliny, 6H-1,2,5-thiadaziny, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 15 thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triaziny, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthy. Preferred heterocyclic ring structures include, but are not limited to, pyridiny, furanyl, thienyl, pyrroly, pyrazolyl, pyrrolidiny, imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazoliny, or isatinoyl. Also included are fused ring 20 and spiro compounds containing, for example, the above heterocyclic ring structures.

As used herein the term "aromatic heterocyclic ring system" has essentially the same definition as for the monocyclic and bicyclic ring systems except that at least one ring of the ring system is an aromatic heterocyclic ring or the bicyclic ring has an aromatic or non-aromatic heterocyclic ring fused to an aromatic carbocyclic ring 25 structure.

The terms "halo" or "halogen" as used herein refer to Cl, Br, F or I substituents. The term "haloalkyl", and the like, refer to an aliphatic carbon radicals having at least one hydrogen atom replaced by a Cl, Br, F or I atom, including mixtures of different halo atoms. Trihaloalkyl includes trifluoromethyl and the like as 30 preferred radicals, for example.

The term "methylene" refers to $-\text{CH}_2-$.

The term "pharmaceutically acceptable salts" includes salts of compounds derived from the combination of a compound and an organic or inorganic acid. These compounds are useful in both free base and salt form. In practice, the use of the salt
5 form amounts to use of the base form; both acid and base addition salts are within the scope of the present invention.

"Pharmaceutically acceptable acid addition salt" refers to salts retaining the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid,
10 hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

15 "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases include salts of primary,
20 secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine,
25 ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine, dicyclohexylamine, choline, and caffeine.


"Biological property" for the purposes herein means an *in vivo* effector or
30 antigenic function or activity that is directly or indirectly performed by a compound of

the invention that are often shown by *in vitro* assays. Effector functions include receptor or ligand binding, any enzyme activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to an extracellular matrix or cell surface molecules, or any structural role. Antigenic functions include possession of an epitope or antigenic site that is capable of reacting with antibodies raised against it.

In the compounds of the invention, carbon atoms bonded to four non-identical substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described herein may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods, or by other methods known in the art. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the asymmetric carbon atoms, when present in the compounds of the invention, may be in one of two configurations (R or S) and both are within the scope of the present invention.


Compounds

The invention provides a compound of following formula I:



wherein:

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$;
 $-C(=O)NR^3R^4$;



- 5 where R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^6 taken
 10 with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

- 15 m is an integer from 0-3, preferably 0-2;

Z is a member selected from the group consisting of a direct link, C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

20

n is an integer from 0-3, preferably 0-2;

- D is a member selected from the group consisting of a direct link, $-CH_2-$, $-O-$, $-N(R^2)-$, $-C(=O)-$, $-S-$, $-SO_2-$, $-SO_2-N(R^2)-$, $-N(R^2)-SO_2-$, $-OC(=O)-$, $-C(=O)O-$,
 25 $-C(=O)-N(R^2)-$ and $-N(R^2)-C(=O)-$, where R^2 is as described above;

R^1 is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -C(=O)NR²R³,
 5 -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, -SO₂R², C_{0-8} alkyl-C(=O)OH and C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, and where R² and R³ are as described above;

10

q is an integer from 0-3, preferably 0-2;

R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl,
 15 C_{1-6} alkyl- C_{3-8} cycloalkyl, -O-R², -O-C(=O)R², - C_{1-8} alkyl-O-R¹⁰, - C_{1-8} alkyl-O-C(=O)R¹⁰, - C_{1-8} alkyl-C(=O)OR¹⁰, - C_{1-8} alkyl-O-C(=O)OR¹⁰, - C_{1-8} alkyl-C(=O)NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, , where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
 20 form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

X is N or -CR¹¹; where R¹¹ is defined as above;


25 p is an integer from 0-3, preferably 0-2;

E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)-, where R¹¹ is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
 30 consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic

ring structure may be independently substituted by from 0 to 5 R^{14} groups;

- Q is a member selected from the group consisting of a direct link, a bivalent C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a
- 5 five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups;
- 10 each R^{14} group is independently a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or
- 15 substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH and C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,


u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are

- 5 independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;
- 10 where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;


with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

- 15 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In a preferred embodiment, the present invention provides a compound of formula I:



- 5 A is a member selected from the group consisting of: R²; -NR³R⁴; -C(=O)NR³R⁴;



- wherein R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are independently selected from the group consisting of H, -OH, C₁₋₆alkyl, C₃₋₈cycloalkyl, C₆₋₁₀aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₄alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R⁶ taken with either of R⁷ and R⁸, and/or R⁷ taken with R⁸, can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

m is an integer from 0-3, preferably 0-2;

Z is a member selected from the group consisting of a direct link, C₁₋₆alkyl,

C_{3-8} cycloalkyl, C_{1-6} alkenyl, C_{6-10} aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

n is an integer from 0-3, preferably 0-2;

D is a member selected from the group consisting of a direct link, $-CH_2-$, $-O-$,
 5 $-NR^2$, $-C(=O)-$, $-S-$, $-SO_2-$, $-SO_2-NR^2$, $-NR^2-SO_2$, $-OC(=O)-$, $-C(=O)NR^2$, and
 $-NR^2-C(=O)-$, where R^2 is as described above;

R^1 is a member selected from the group consisting of H, C_{1-6} alkyl, halogen,
 $-C(=O)OH$, an unsubstituted amino group, a mono- or di-substituted amino group,
 10 $-CN$, $-NO_2$, $-OH$, $-C(=O)NR^2R^3$, $-OR^2$ and $-O-C(=O)R^2$, and where R^2 and R^3 are as
 described above;

q is an integer from 0-3, preferably 0-2;

R^{11} is a member selected from the group consisting of H, C_{1-6} alkyl,
 15 C_{3-8} cycloalkyl, C_{6-10} aryl, C_{1-4} alkylaryl, C_{1-4} alkyl- C_{3-8} cycloalkyl, $-OR^2$, $-O-C(=O)R^2$,
 $-C_{1-6}$ alkyl- $O-R^{10}$, $-C_{1-6}$ alkyl- $O-C(=O)R^{10}$, $-C_{1-6}$ alkyl- $C(=O)OR^{10}$,
 $-C_{1-6}$ alkyl- $O-C(=O)OR^{10}$, $-C_{1-6}$ alkyl- $C(=O)NR^{10}R^{10}$, $-C_{1-6}$ alkyl- $NR^{10}R^{10}$,
 $-C_{1-6}$ alkyl- $NR^{10}C(=O)R^{10}$, $-SR^{10}$, where R^2 is as described above and R^{10} is a member
 20 selected from the group consisting of H, C_{1-6} alkyl, and wherein when two R^{10} groups
 are present they may be taken together to form a saturated or unsaturated ring with the
 atom to which they are both attached;

X is N or $-CR^{11}$; where R^{11} is defined as above;


p is an integer from 0-3, preferably 0-2;

E is a member selected from the group consisting of a direct link, $-O-$, $-NR^{11}$,
 25 where R^{11} is as defined above, phenyl, a 5 to 12 member bivalent heteroaryl group
 containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a
 five to ten membered non-aromatic heterocyclic ring system containing 1-4
 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl
 and said non-aromatic heterocyclic ring structure may be independently substituted by

from 0 to 5 R^{14} groups and each R^{14} group is independently defined as set forth above for R^1 ;

- Q is a member selected from the group consisting of a direct link, C_{3-8} cycloalkyl, phenyl, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups and each R^{14} group is independently defined as set forth above for R^1 ;

- 10 G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,

- u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} ,


can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

- 5 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In another preferred aspect the present invention provides a compound of formula I as illustrated above wherein:

- 10 A is a member selected from the group consisting of: R²; -NR³R⁴; -C(=O)NR³R⁴;



- where R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R⁶ taken with either of R⁷ and R⁸, and/or R⁷ taken with R⁸, can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;
- 20 m is 0;

Z is a member selected from the group consisting of a direct link, C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₁₋₈carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

5

n is 0;

D is a member selected from the group consisting of a direct link, -CH₂-, -O-, -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-,
10 -C(=O)-N(R²)- and -N(R²)-C(=O)-, where R² is as described above;

R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³,
15 -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² and R³ are as described above;

20

q is an integer from 0-3, preferably 0-2;

R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl,
25 C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
30 form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;


X is N;

p is 0-2;

- 5 E is a member selected from the group consisting of a direct link, -O-,
-N(-R¹¹)-, where R¹¹ is as defined above, phenylene, a bivalent 5 to 12 member
bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
consisting of N, O and S, and a five to ten membered non-aromatic bivalent
heterocyclic ring system containing 1-4 heteroatoms selected from the group
10 consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic
ring structure may be independently substituted by from 0 to 5 R¹⁴ groups and each
R¹⁴ group is independently defined the same as the substituents set forth above for the
R¹ group;

- Q is a member selected from the group consisting of a direct link, a bivalent
15 C₃₋₈cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group
containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a
five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4
heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl
and said non-aromatic heterocyclic ring structure may be independently substituted by
20 from 0 to 5 R¹⁴ groups and each R¹⁴ group is independently defined the same as the
substituents set forth above for the R¹ group;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,

5 u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;


10 where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at

15 least one N atom;


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

Particularly preferred benzimidazole compounds of the invention are compounds of formula II:



wherein:

- 5 A is a member selected from the group consisting of: R²; -NR³R⁴; -C(=O)NR³R⁴;



- where R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are independently selected from the group
- 10 consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R⁶ taken
- 15 with either of R⁷ and R⁸, and/or R⁷ taken with R⁸, can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

m is 0;

Z is a member selected from the group consisting of a direct link, C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₁₋₈carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

5

n is 0;

D is a member selected from the group consisting of: -CH₂-, -O-, -NR², -C(=O)-, -S-, -SO₂-, -SO₂-NR², -NR²-SO₂-, -OC(=O)-, -C(=O)NR², and -NR²-C(=O)-, and is preferably a member selected from the group consisting of: -O-, -NR², -C(=O)-, -S-, -SO₂-, where R² is as set forth above;

10

q is an integer from 0-3;

15

R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

20

25

p is an integer from 0-2;


E is a member selected from the group consisting of a direct link, -O-, -N(-R¹¹)-, where R¹¹ is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴ groups and each

30

R¹⁴ group is independently defined the same as the substituents set forth above for the R¹ group;

Q is a member selected from the group consisting of a direct link, a bivalent C₃₋₈cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴ groups and each R¹⁴ group is independently defined the same as the substituents set forth above for the R¹ group;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,


u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

Further preferred are compounds of formula III:



wherein:

R^2 and R^8 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

R^1 is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, -SO₂R², C_{0-8} alkyl-C(=O)OH and

C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, and where R^2 and R^3 are as described above;

q is an integer from 0-3;

5 R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl, C_{1-6} alkyl- C_{3-8} cycloalkyl, -O- R^2 , -O-C(=O) R^2 , - C_{1-8} alkyl-O- R^{10} , - C_{1-8} alkyl-O-C(=O) R^{10} , - C_{1-8} alkyl-C(=O)OR¹⁰, - C_{1-8} alkyl-O-C(=O)OR¹⁰, - C_{1-8} alkyl-C(=O)NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰C(=O) R^{10} , -SR¹⁰, where R^2 is as described above

10 and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;


15 p is an integer from 0-2;

E is a member selected from the group consisting of a direct link, -O-, -N(- R^{11})-, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
 20 consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups and each R^{14} group is independently defined the same as the substituents set forth above for the
 25 R^1 group;

Q is a member selected from the group consisting of a direct link, a bivalent C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4
 30 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl

and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups and each R^{14} group is independently defined the same as the substituents set forth above for the R^1 group;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



5 wherein

t is an integer from 0 to 6,

u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;


with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

Particularly preferred compounds of formula III are compounds wherein R¹ and R⁸ are each independently a lower alkyl group and R¹¹ is C₃-C₈ cycloalkyl group. Particularly preferred cycloalkyl groups are members selected from the group consisting of cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. Further preferred compounds of formula III are such compounds wherein one or more of E and Q is independently a phenylene, a heteroaryl or a heterocyclic group as defined above with respect to formula II, especially a phenylene, a heteroaryl or heterocyclic member selected from the group consisting of phenyl, thiophene, furan, benzofuran, benzothiophene and pyridine. Other heterocyclic bicyclic rings as defined above for formula II, and the like may be used for E and Q as well. When only one of E or Q is a phenylene, heteroaryl or heterocyclic member, the other is preferably a direct link. More preferred are compounds wherein q is zero and R⁸ is lower alkyl.

Even more preferred compounds of formula III are such compounds according to formula IIIa as set forth in Table 1, below.


Table 1



Formula IIIa

p	E	Q	G
1	direct link		
1	direct link		
1	direct link		
1	direct link		
1	direct link		
2	direct link		
1			
1			
2			


Table 1 (Cont.)



Formula IIIa

p	E	Q	G
1			
1			
1			
1			
1			
1			

Also preferred compounds are benzimidazole compounds according to formula II wherein D is -O-, p and q are each an integer from 1-3, and E and Q collectively form a substituted or unsubstituted indole group. More preferred compounds having such an indole group are compounds according to formula IV as set forth below (and having the same defined remaining substituents as in formula II, above).




(Formula IV)





10 Particularly preferred compounds according to formula IV are compounds wherein each of the R¹, R⁸ and R¹⁴ groups is independently selected from the group consisting of hydrogen and C₁-C₃ alkyl, preferably, hydrogen and C₁-C₃ alkyl, and most preferably, methyl and ethyl groups.

Even more more preferred compounds according to formula IV are
15 compounds according to formula IVa as set forth in Table 2, below.

Table 2




Formula IVa

R ¹	R ¹¹
-H	-CH ₃
-OH	-CH ₂ -CH ₃
-NH ₂	-CH(-CH ₃) ₂
-NHCH ₂ -COOH	-CH ₂ COOH
-NHSO ₂ -CH ₃	-CH ₂ - 
-Br	
-COOH	
-COOCH ₃	
-CF ₃	-CH ₂ CH ₂ OH
-CONH ₂	-C(-CH ₃) ₃


Also preferred compounds are benzimidazole compounds according to formula II wherein m is zero and E and Q collectively form a substituted or unsubstituted biphenylene group. More preferred compounds having such an biphenylene group are compounds according to formula V as set forth below.

5



wherein:

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$; $-C(=O)NR^3R^4$;



10

where R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and

15 C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered

heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

5 Z is a member selected from the group consisting of a direct link, C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₁₋₈carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

n is an integer from 0-3, preferably 0-2;

10

D is a member selected from the group consisting of a direct link, -CH₂-, -O-, -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-, -C(=O)-N(R²)- and -N(R²)-C(=O)-, where R² is as described above;

15

R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² and R³ are as described above;

20

q is an integer from 0-3, preferably 0-2;

25


R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to

30

form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

- each R^{14} group is independently a member selected from the group consisting
- 5 of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$, C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl, -CN, $-NO_2$, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, $-O-R^2$ and $-O-C(=O)R^2$, an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl,
- 10 C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$ and C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl;

G is a member selected from the group consisting of:



wherein

- t is an integer from 0 to 6,
- 15 u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms
- 20 with 1 to 4 of such atoms being selected from the group consisting of N, O and S;


where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

- and all pharmaceutically acceptable isomers, salts, hydrates, solvates and
5 prodrug derivatives thereof.

Particularly preferred compounds according to formula V are compounds wherein each of the R^1 and R^{14} groups is independently selected from the group consisting of hydrogen and C_1 - C_5 alkyl, preferably, hydrogen and C_1 - C_3 alkyl, most preferably, hydrogen, methyl and ethyl; and R^{11} is H, or C_3 to C_8 cycloalkyl.

- 10 Even more more preferred compounds according to formula V are compounds according to formula Va as set forth in Table 3, below.

Table 3




Formula Va

5

A	Z	n	D
		0	O
		0	
		0	O
		0	O
CH ₃ -		2	O
H-		2	O
NH ₂ -		2	O

Table 3 (Cont.)




Formula Va

A	Z	n	D
H-		2	O
CH ₃ -		2	O
(CH ₃) ₂ NCO-		0	
		0	O
		1	CH ₂

5

10 Also preferred compounds are compounds according to formula II having two bicyclic ring structures as set forth in the definition for formula II. Particularly preferred such compounds are compounds wherein at least one of the bicyclic rings is

joined directly or indirectly to a piperidine ring wherein D is -O- and each of the remaining substituents is defined as set forth in formula II as illustrated by formula VI as set forth below:



5

Formula VI

wherein:

R^6 and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten
10 membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

15 R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl, C_{1-6} alkyl- C_{3-8} cycloalkyl, -O- R^2 , -O-C(=O) R^2 , - C_{1-8} alkyl-O- R^{10} , - C_{1-8} alkyl-O-C(=O) R^{10} , - C_{1-8} alkyl-C(=O)OR¹⁰, - C_{1-8} alkyl-O-C(=O)OR¹⁰, - C_{1-8} alkyl-C(=O)NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R^2 is as described above
20 and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

X is N or $-CR^{11}$; where R^{11} is defined as above;


p is an integer from 0-3;

E is a member selected from the group consisting of a direct link, -O-,
5 -N($-R^{11}$)-, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member
bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
consisting of N, O and S, and a five to ten membered non-aromatic bivalent
heterocyclic ring system containing 1-4 heteroatoms selected from the group
consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic
10 ring structure may be independently substituted by from 0 to 5 R^{14} groups;

each R^{14} group is independently a member selected from the group consisting
of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl,
 C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH,
15 C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or
di-substituted amino group, wherein the substituted amino groups are independently
substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl,
 C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH and
 C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl;

20 Q is a member selected from the group consisting of a direct link, a bivalent
 C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group
containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a
five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4
heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl
25 and said non-aromatic heterocyclic ring structure may be independently substituted by
from 0 to 5 R^{14} groups;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,


u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are

- 5 independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;
- 10 where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

- 15 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

More preferred compounds are those wherein E and Q collectively form a substituted or unsubstituted indole group as set forth in formula VIa, below:



wherein:

R^6 and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten
 5 membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl,
 10 C_{1-6} alkyl- C_{3-8} cycloalkyl, -O- R^2 , -O-C(=O) R^2 , - C_{1-8} alkyl-O- R^{10} , - C_{1-8} alkyl-C(=O) R^{10} , - C_{1-8} alkyl-O-C(=O) R^{10} , - C_{1-8} alkyl-O-C(=O) R^{10} , - C_{1-8} alkyl-C(=O) $NR^{10}R^{10}$, - C_{1-8} alkyl- $NR^{10}R^{10}$, - C_{1-8} alkyl- $NR^{10}C(=O)R^{10}$, -SR¹⁰, where R^2 is as described above
 15 and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached, in which case preferably form a partially or fully saturated ring;

20 X is N or -CR¹¹; where R^{11} is defined as above;

p is an integer from 0-3;

each R^{14} group is independently a member selected from the group consisting
 25 of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or

di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

5

R²³, R²⁴ and R²⁵ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the

10 ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


15

Even more preferred are such compounds according to formula VIa wherein p is an integer from 1-3, R⁶ is hydrogen, R⁹ is C₁-C₃ alkyl and the remaining substituents are defined as set forth in the definitions of formula II, above.

20

More preferred compounds are compounds having such G group, wherein R²³ is H or C₁-C₅ alkyl and R²⁴ and R²⁵ are independently selected from H and C₁-C₅ alkyl.

Especially preferred such compounds are compounds according to formula VIb as set forth below (and having the same defined remaining substituents as in formula II, above):




5

(Formula VIb)

Examples of such particularly preferred compounds according to formula VIb are set forth below in Table 4.


Table 4



Formula VIb

X	p	R ⁹	R ¹¹	R ¹⁴
N	1	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
N	2	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
N	3	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
N	0	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
N	1	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
N	2	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
N	3	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
N	0	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
N	1	-CH ₃	-COOCH ₂ CH ₃	-CH ₃
N	1	-CH ₃	-CH(CH ₃) ₂	-CH ₃

Table 4 (Cont.)






Formula VIb

X	P	R ⁹	R ¹¹	R ¹⁴
C	1	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
C	2	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
C	3	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
C	0	-CH ₃	-COOCH ₂ CH ₃	-CH ₂ CH ₃
C	1	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
C	2	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
C	3	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
C	0	-CH ₃	-CH(CH ₃) ₂	-CH ₂ CH ₃
C	1	-CH ₃	-COOCH ₂ CH ₃	-CH ₃
C	1	-CH ₃	-CH(CH ₃) ₂	-CH ₃

Some preferred embodiments of the invention are shown in the following Table 5.

Table 5

5	STRUCTURE	Inhibitory Activity (IC ₅₀)	
		Factor Xa	Thrombin
		1.2 nM	805 nM
		18 nM	1.3 μM
		5.73 nM	26.3 μM

The invention also encompasses all pharmaceutically acceptable salts, hydrates, solvates, and prodrug derivatives of the compounds of formulae I-VIb. In addition, the compounds of formulae I-VIb can exist in various isomeric and tautomeric forms, and all such forms are meant to be included in the invention, along

with pharmaceutically acceptable salts, hydrates, solvates, and prodrug derivatives of such isomers and tautomers.

The compounds of the invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of the invention. Non-toxic and physiologically compatible salts are particularly useful, but less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, the free acid or free base form of a compound of one of the formulas above can be reacted with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

Prodrug Derivatives of Compounds

The invention also encompasses prodrug derivatives of the compounds contained herein. The term "prodrug" refers to a pharmacologically inactive derivative of a parent drug molecule that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active drug. Prodrugs are variations or derivatives of the compounds of the invention which have groups cleavable under metabolic conditions. Prodrugs become the compounds of the invention which are pharmaceutically active *in vivo*, when they undergo solvolysis under physiological conditions or undergo enzymatic degradation. Prodrug compounds of the invention may be called single, double, triple etc., depending on the number of biotransformation steps required to release the active drug within the organism, and indicating the number of functionalities present in a precursor-type form. Prodrug forms often offer advantages of solubility, tissue compatibility, or

delayed release in the mammalian organism (see, Bundgard, Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985 and Silverman, The Organic Chemistry of Drug Design and Drug Action, pp. 352-401, Academic Press, San Diego, CA, 1992). Prodrugs commonly known in the art include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acids with a suitable alcohol, or amides prepared by reaction of the parent acid compound with an amine, or basic groups reacted to form an acylated base derivative. Moreover, the prodrug derivatives of the invention may be combined with other features herein taught to enhance bioavailability.

10 The compounds of the present invention may also be used alone or in combination or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of the invention may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen
15 activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of the present invention may act in a synergistic fashion to prevent reocclusion following a successful thrombolytic therapy and/or reduce the time to reperfusion. These compounds may also allow for reduced doses of the thrombolytic
20 agents to be used and therefore minimize potential hemorrhagic side-effects. The compounds of the invention can be utilized *in vivo*, ordinarily in mammals such as primates, (e.g. humans), sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

25 The biological properties of the compounds of the present invention can be readily characterized by methods that are well known in the art such as, for example, by the *in vitro* protease activity assays and *in vivo* studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters, such as are illustrated in the examples.

30 Diagnostic applications of the compounds of the invention will typically utilize formulations in the form of solutions or suspensions. In the management of

thrombotic disorders, the compounds of the invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in need of treatment (typically mammalian) using the compounds of the invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these compounds are employed, and other factors which those skilled in the medical arts will recognize.

Preparation of Compounds

The compounds of the present invention may be synthesized by either solid or liquid phase methods described and referenced in standard textbooks, or by a combination of both methods. These methods are well known in the art. See, Bodanszky, "The Principles of Peptide Synthesis", Hafner, *et al.*, Eds., Springer-Verlag, Berlin, 1984.


Starting materials used in any of these methods are commercially available from chemical vendors such as Aldrich, Sigma, Nova Biochemicals, Bachem Biosciences, and the like, or may be readily synthesized by known procedures.

Reactions are carried out in standard laboratory glassware and reaction vessels under reaction conditions of standard temperature and pressure, except where otherwise indicated.


During the synthesis of these compounds, the functional groups of the amino acid derivatives used in these methods are protected by blocking groups to prevent cross reaction during the coupling procedure. Examples of suitable blocking groups and their use are described in "The Peptides: Analysis, Synthesis, Biology", Academic Press, Vol. 3 (Gross, *et al.*, Eds., 1981) and Vol. 9 (1987), the disclosures of which are incorporated herein by reference.

One exemplary synthesis scheme is outlined directly below, and the specific steps are described in the Examples. The reaction products are isolated and purified by conventional methods, typically by solvent extraction into a compatible solvent. The products may be further purified by column chromatography or other appropriate methods.

Scheme 1



Scheme 2



Compositions or Formulations

Compositions or formulations of the compounds of the invention are prepared for storage or administration by mixing a compound of the invention having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as TWEEN®, PLURONICS® or polyethyleneglycol.

Dosage formulations of the compounds of the invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of the invention typically will be about 3-11, more preferably about 5-9 and most preferably about 7-8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers may result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as orally, intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally, transdermally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such as ointments, drops and

dermal patches. The compounds of the invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

5 The compounds of the invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.

 The compounds of the invention may also be delivered by the use of
10 antibodies, antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of the invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidinone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-
15 polylysine substituted with palmitoyl residues. Furthermore, compounds of the invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans,
20 polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

 Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial
25 having a stopper pierceable by hypodermic injection needle.

 Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will be influenced by the route of administration, the
30 therapeutic objectives and the condition of the patient. For injection by hypodermic

needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each compound by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be readily determined by one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

The compounds and compositions/formulations of the invention can be administered orally or parenterally in an effective amount within the dosage range of about 0.001 to about 1000 mg/kg, preferably about 0.01 to about 100 mg/kg and more preferably about 0.1 to about 20 mg/kg. Advantageously, the compounds and compositions/formulations of the invention may be administered several times daily, although other dosage regimens may also be useful (e.g. single daily dose and/or continuous infusion).

Typically, about 0.5 to about 500 mg of at least one compound or mixture of compounds of the invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are binders such as acacia, corn starch or gelatin, and excipients such as microcrystalline cellulose, disintegrating agents like corn starch or alginic acid, lubricants such as magnesium stearate, sweetening agents such as sucrose or lactose, or flavoring agents. When a dosage form is a capsule, in addition to the above materials it may also contain liquid carriers such as water, saline, or a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated

according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

The preferred compounds of the present invention are characterized by their ability to inhibit thrombus formation with acceptable effects on classical measures of coagulation parameters, platelets and platelet function, and acceptable levels of bleeding complications associated with their use. Conditions characterized by undesired thrombosis would include those involving the arterial and venous vasculature.

With respect to the coronary arterial vasculature, abnormal thrombus formation characterizes the rupture of an established atherosclerotic plaque which is the major cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA).

With respect to the venous vasculature, abnormal thrombus formation characterizes the condition observed in patients undergoing major surgery in the lower extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombus formation further characterizes disseminated intravascular coagulopathy commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.

The compounds of the invention are useful for the treatment or prophylaxis of those diseases which involve the production and/or action of factor


Xa/prothrombinase complex. The compounds of this present invention, selected and used as disclosed herein, find utility as a diagnostic or therapeutic agent for preventing or treating a condition in a mammal characterized by undesired thrombosis or a disorder of coagulation. Disease states treatable or preventable by the administration of compounds of the invention include, without limitation, occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty, thrombus formation in the venous vasculature, disseminated intravascular coagulopathy, the treatment of reocclusion or restenosis of reperfused coronary arteries, thromboembolic complications of surgery and peripheral arterial occlusion, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure, hemorrhagic stroke, renal dialysis, blood oxygenation, and cardiac catheterization.

Accordingly, the invention provides a method for preventing or treating a condition in a mammal characterized by undesired thrombosis which administers to a mammal a therapeutically effective amount of a compound of the invention, as described herein. Conditions for prevention or treatment include, for example, (a) the treatment or prevention of any thrombotically mediated acute coronary syndrome including myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, (b) the treatment or prevention of any thrombotically mediated cerebrovascular syndrome including embolic stroke, thrombotic stroke or transient ischemic attacks, (c) the treatment or prevention of any thrombotic syndrome occurring in the venous system including deep venous thrombosis or pulmonary embolus occurring either spontaneously or in the setting of malignancy, surgery or trauma, (d) the treatment or prevention of any coagulopathy including disseminated intravascular coagulation (including the setting of septic shock or other infection, surgery, pregnancy, trauma or malignancy and whether associated with multi-organ failure or not), thrombotic thrombocytopenic purpura, thromboangiitis obliterans, or thrombotic disease associated with heparin induced thrombocytopenia, (e) the treatment or prevention of thrombotic complications associated with extracorporeal circulation (e.g. renal


dialysis, cardiopulmonary bypass or other oxygenation procedure, plasmapheresis), (f) the treatment or prevention of thrombotic complications associated with instrumentation (e.g. cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve), and (g) those involved with the fitting of
5 prosthetic devices.

Anticoagulant therapy is also useful to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus the compounds of the invention can be added to or contacted with any medium
10 containing or suspected to contain factor Xa and in which it is desired that blood coagulation be inhibited, e.g., when contacting the mammal's blood with material such as vascular grafts, stents, orthopedic prostheses, cardiac stents, valves and prostheses, extra corporeal circulation systems and the like. Thus, the compounds of the invention also find utility in a method for inhibiting the coagulation of biological
15 samples by administration of a compound of the invention.


Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and
20 utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

EXAMPLESExample 1


- 5 To a solution of tert-Butylamine (41.4g, 566 mmol) and triethylamine (118 mL, 849 mmol) in DCM (1000 mL) in an ice bath, was added benzenesulfonyl chloride (100 g, 566 mmol) dropwise. The mixture was stirred at room temperature overnight. Water was added to the mixture and organic layer was washed with water, brine, dried over Na₂SO₄, filtered and filtrate evaporated *in vacuo* to give the title compound as
- 10 light yellowish solid (117.63 g, 97.6%). ES-MS (M+H)⁺ = 214.5.

Example 2


- To a solution of compound of example 1 (53.25 g, 250 mmol) in THF (600 mL)
- 15 in an ice bath, was added n-butyllithium in hexane (200 mL, 500 mmol) dropwise. A thick precipitate was formed when the reaction mixture was warmed up to 10°C. Triisopropylborate was added keeping the temperature below 35°C. After 1 hr., the mixture was cooled in an ice bath, 1N HCl (405 mL) was added, and the mixture was stirred overnight. The mixture was extracted with ether (100 mL) three times. The
- 20 combined organic extracts were extracted with 1N NaOH (130 mL) three times. The aqueous extracts were acidified to pH 1 with 12 N HCl, and then extracted with ether three times (140 ML). The combined ether extracts were dried over MgSO₄, and solvents evaporated in vacuo. Hexane and ether were added and a white precipitate formed. The solid was collected and washed with 10% ether/hexane to give the title
- 25 compound. ES-MS (M+H)⁺ = 257.8.

Example 3

To a solution of 5-bromoindole (1.96 g, 10 mmol) in DME (40 mL) and H₂O (10 mL), was added compound of example 2 (3.85 g, 15 mmol), NaHCO₃ (1.68g, 20 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.58g, 0.5 mmol). The mixture was refluxed overnight, cooled to room temperature, diluted with EtOAc. The organic layer was washed with water, dried with MgSO₄, filtered and concentrated. This was purified by silica gel column chromatography using solvent system 25% EtOAc in hexane as eluant to give the title compound (1.52g, 46%). ES-MS (M+Na)⁺ = 351.1.

Example 4

To a solution of compound of example 3 (328 mg, 1 mmol) in DMF (10 mL) was added (296 mg, 1.2 mmol) and Cs₂CO₃ (1.3g, 4 mmol). The mixture was stirred at room temperature overnight. Water and EtOAc were added. The organic layer was washed with water, 1N HCl, brine, dried over Na₂SO₄ and concentrated *in vacuo*. It was purified by silica gel column chromatography using solvent system 50% EtOAc in DCM followed by DCM as eluant to give the title compound (237 mg, 48%). ES-MS (M+H)⁺ = 494.2.


Example 5

- A solution of the compound of example 4 (115mg, 0.23 mmol) in 20% MeOH/EtOAc (10 mL) was treated with a stream of HCl gas for 10 min. at 0°C. The resulting solution was capped, stirred at room temperature overnight and evaporated *in vacuo*. The residue was reconstituted in MeOH (10 mL) and the mixture was treated with NH₄OAc (350 mg, 4.6 mmol). The reaction mixture was refluxed for 2 hrs. and concentrated *in vacuo*. The obtained residue was purified by RP-HPLC to give the title compound as a white powder. ES-MS (M+H)⁺ = 511.2.


Further compound Examples 6-21 were made by adapting the procedures of Examples 1-5, to produce compounds according to the following formulae:

Example






6







7



Example




Example




Example


17




18




19



20



21



Below is biological data reported as IC₅₀ values, as described below, for compound Examples 6-18, respectively, as shown above.

Example	XA	IIA	IIASE	TRYPSIN
6	0.148	0.0164	0.139	1.66
7	0.0012	0.805	0.00967	0.167
8	0.0322	29	0.375	6.13
9	0.658	9.61	9.04	4.92
10	0.00573	26.3	0.00256	0.832
11	0.0104		0.017	
12	0.0584		0.124	
13	0.386	102	8.1	0.711
14	0.0165	52.5	0.182	0.176
15	0.031	23.1	0.277	0.105
16	0.0484	330	0.45	0.077
17	0.0199	305	0.38	0.0725
18	0.018	1.357	0.167	0.109

5

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description, make and utilize the compounds of the present invention and practice the claimed methods.

10

BIOLOGICAL ACTIVITY EXAMPLES

Evaluation of the compounds of the invention is guided by in vitro protease activity assays (see below) and in vivo studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters.

15

The compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from about 0

to about 100 μ M. In the assays for thrombin, prothrombinase and factor Xa, a synthetic chromogenic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The IC_{50} of a compound is determined from the substrate turnover. The IC_{50} is the concentration of test compound giving about 50% inhibition of the substrate turnover. The compounds of the present invention desirably have an IC_{50} of less than about 500 nM in the factor Xa assay, preferably less than about 200 nM, and more preferred compounds have an IC_{50} of about 100 nM or less in the factor Xa assay. The compounds of the present invention desirably have an IC_{50} of less than about 4.0 μ M in the prothrombinase assay, preferably less than about 200 nM, and more preferred compounds have an IC_{50} of about 10 nM or less in the prothrombinase assay. The compounds of the present invention desirably have an IC_{50} of greater than about 1.0 μ M in the thrombin assay, preferably greater than about 10.0 μ M, and more preferred compounds have an IC_{50} of greater than about 100.0 μ M in the thrombin assay.

Amidolytic Assays for determining protease inhibition activity

The factor Xa and thrombin assays were performed at room temperature, in 0.02 M Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl. The rates of hydrolysis of the para-nitroanilide substrate S-2765 (Chromogenix) for factor Xa, and the substrate Chromozym TH (Boehringer Mannheim) for thrombin following preincubation of the enzyme with inhibitor for 5 minutes at room temperature, and were determined using the Softmax 96-well plate reader (Molecular Devices), monitored at 405 nm to measure the time dependent appearance of p-nitroaniline.

25

The prothrombinase inhibition assay was performed in a plasma free system with modifications to the method described by Sinha, U. *et al.*, *Thromb. Res.*, 75, 427-436 (1994). Specifically, the activity of the prothrombinase complex was determined by measuring the time course of thrombin generation using the p-nitroanilide substrate Chromozym TH. The assay consists of preincubation (5 minutes) of selected compounds to be tested as inhibitors with the complex formed

30

from factor Xa (0.5 nM), factor Va (2 nM), phosphatidyl serine:phosphatidyl choline (25:75, 20 μ M) in 20 mM Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl, 5 mM CaCl₂ and 0.1% bovine serum albumin. Aliquots from the complex-inhibitor mixture were added to prothrombin (1 nM) and Chromozym TH (0.1 mM). The rate of substrate cleavage was monitored at 405 nm for two minutes. Eight different concentrations of inhibitor were assayed in duplicate. A standard curve of thrombin generation by an equivalent amount of untreated complex was used for determination of percent inhibition.

10 A series of studies are accomplished in rabbits to evaluate the antithrombotic efficacy, and effects on hemostasis and hematological parameters of the above compounds as follows.

Antithrombotic Efficacy in a Rabbit Model of Venous Thrombosis

15 A rabbit deep vein thrombosis model as described by Hollenbach, S. et al., Thromb. Haemost. 71, 357-362 (1994), is used to determine the in-vivo antithrombotic activity of the test compounds. Rabbits are anesthetized with I.M. injections of Ketamine, Xylazine, and Acepromazine cocktail. A standardized protocol consists of insertion of a thrombogenic cotton thread and copper wire apparatus into the abdominal vena cava of the anesthetized rabbit. A non-occlusive thrombus is allowed to develop in the central venous circulation and inhibition of thrombus growth is used as a measure of the antithrombotic activity of the studied compounds. Test agents or control saline are administered through a marginal ear vein catheter. A femoral vein catheter is used for blood sampling prior to and during steady state infusion of test compound. Initiation of thrombus formation begins immediately after advancement of the cotton thread apparatus into the central venous circulation. Test compounds are administered from time = 30 min to time = 150 min at which the experiment is terminated. The rabbits are euthanized and the thrombus excised by surgical dissection and characterized by weight and histology. Blood samples are analyzed for changes in hematological and coagulation parameters.

Effects of Compounds of the Invention in Rabbit Venous Thrombosis model


Administration of above compounds in the rabbit venous thrombosis model demonstrates antithrombotic efficacy at the higher doses evaluated. There are no significant effects of the compound on the aPTT and PT prolongation with the highest dose (100 $\mu\text{g/kg}$ + 2.57 $\mu\text{g/kg/min}$). Compound xxxx has no significant effects on hematological parameters as compared to saline controls. All measurements are an average of all samples after steady state administration of vehicle or (D)-Arg-Gly-Arg-thiazole. Values are expressed as mean \pm SD.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as follows in the scope of the appended claims.

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference.

WHAT IS CLAIMED IS:

1. A compound represented by the formula:




5

wherein:

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$;
 $-C(=O)NR^3R^4$;

10



15

wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of

N, O and S;

m is an integer from 0-3;

- 5 Z is a member selected from the group consisting of a direct link, C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₁₋₈carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

n is an integer from 0-3;

10

D is a member selected from the group consisting of a direct link, -CH₂-, -O-, -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-, -C(=O)-N(R²)- and -N(R²)-C(=O)-, where R² is as described above;

15

R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² and R³ are as described above;

20

q is an integer from 0-3;

25

- R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to
- 30

form a saturated or unsaturated ring with the atom to which they are both attached;

X is N or $-CR^{11}$; where R^{11} is defined as above;


5 p is an integer from 0-3;

E is a member selected from the group consisting of a direct link, -O-,
-N(R^{11})-, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member
bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
10 consisting of N, O and S, and a five to ten membered non-aromatic bivalent
heterocyclic ring system containing 1-4 heteroatoms selected from the group
consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic
ring structure may be independently substituted by from 0 to 5 R^{14} groups;

Q is a member selected from the group consisting of a direct link, a bivalent
15 C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group
containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a
five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4
heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl
and said non-aromatic heterocyclic ring structure may be independently substituted by
20 from 0 to 5 R^{14} groups;

each R^{14} group is independently a member selected from the group consisting
of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl,
 C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH,
 C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or
25 di-substituted amino group, wherein the substituted amino groups are independently
substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl,
 C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH and
 C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,


- u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are
- 5 independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;
- 10 where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

- 15 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

2. A compound of claim 1, wherein:

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$;
 $-C(=O)NR^3R^4$;



- 5 wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-6} alkyl, C_{3-8} cycloalkyl, C_{6-10} aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-4} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group
- 10 consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

m is an integer from 0-3;

- Z is a member selected from the group consisting of a direct link, C_{1-6} alkyl,
- 15 C_{3-8} cycloalkyl, C_{1-6} alkenyl, C_{6-10} aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

n is an integer from 0-3;

- D is a member selected from the group consisting of a direct link, $-CH_2-$, $-O-$, $-NR^2$, $-C(=O)-$, $-S-$, $-SO_2-$, $-SO_2-NR^2$, $-NR^2-SO_2$, $-OC(=O)-$, $-C(=O)NR^2$, and
- 20 $-NR^2-C(=O)-$, where R^2 is as described above;

R^1 is a member selected from the group consisting of H, C_{1-6} alkyl, halogen, $-C(=O)OH$, an unsubstituted amino group, a mono- or di-substituted amino group,

-CN, -NO₂, -OH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², and where R² is as described above;

q is 0-3;

- 5 R¹¹ is a member selected from the group consisting of H, C₁₋₆alkyl, C₃₋₈cycloalkyl, C₆₋₁₀aryl, C₁₋₄alkylaryl, C₁₋₄alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₆alkyl-O-R¹⁰, -C₁₋₆alkyl-O-C(=O)R¹⁰, -C₁₋₆alkyl-C(=O)OR¹⁰, -C₁₋₆alkyl-O-C(=O)OR¹⁰, -C₁₋₆alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₆alkyl-NR¹⁰R¹⁰, -C₁₋₆alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member
- 10 selected from the group consisting of H, C₁₋₆alkyl, and wherein when two R¹⁰ groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached;

X is N or -CR¹¹; where R¹¹ is defined as above;

p is an integer from 0-3;


- 15 E is a member selected from the group consisting of a direct link, -O-, -NR¹¹, where R¹¹ is as defined above, phenyl, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl
- 20 and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴ groups;

- Q is a member selected from the group consisting of a direct link, C₃₋₈cycloalkyl, phenyl, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten
- 25 membered non-aromatic heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴ groups;

each R^{14} group is independently a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or

5 di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH and C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



10 wherein

t is an integer from 0 to 6,

u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic

15 ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1


20 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


5

3. A compound of formula II:



wherein:

10 A is a member selected from the group consisting of: R²; -NR³R⁴; -C(=O)NR³R⁴;



15 where R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and R⁹ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R⁶ taken
20 with either of R⁷ and R⁸, and/or R⁷ taken with R⁸, can each form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of

N, O and S;

m is 0;

- 5 Z is a member selected from the group consisting of a direct link, C_{1-8} alkyl, C_{3-8} cycloalkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{1-8} carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

n is 0;

10

D is a member selected from the group consisting of: $-CH_2-$, $-O-$, $-NR^2$, $-C(=O)-$, $-S-$, $-SO_2-$, $-SO_2-NR^2$, $-NR^2-SO_2$, $-OC(=O)-$, $-C(=O)NR^2$, and $-NR^2-C(=O)-$, where R^2 is as described above;

15

q is an integer from 0-3;

- R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl, C_{1-6} alkyl- C_{3-8} cycloalkyl, $-O-R^2$, $-O-C(=O)R^2$, $-C_{1-8}$ alkyl- $O-R^{10}$, $-C_{1-8}$ alkyl- $O-C(=O)R^{10}$, $-C_{1-8}$ alkyl- $C(=O)OR^{10}$, $-C_{1-8}$ alkyl- $O-C(=O)OR^{10}$, $-C_{1-8}$ alkyl- $C(=O)NR^{10}R^{10}$, $-C_{1-8}$ alkyl- $NR^{10}R^{10}$, $-C_{1-8}$ alkyl- $NR^{10}C(=O)R^{10}$, $-SR^{10}$, where R^2 is as described above and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached;

25

X is N or $-CR^{11}$; where R^{11} is defined as above;

p is an integer from 0-2;


- 30 E is a member selected from the group consisting of a direct link, $-O-$, $-N(-R^{11})-$, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group

consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R¹⁴ groups;

- 5 Q is a member selected from the group consisting of a direct link, a bivalent C₃₋₈cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by
- 10 from 0 to 5 R¹⁴ groups;

- each R¹⁴ group is independently a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH,
- 15 C₀₋₈alkyl-SH, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

5 t is an integer from 0 to 6;


u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic
 10 ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;
 where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵,
 15 can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

4. A compound of formula III:

5



wherein:

10 R^2 and R^8 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

15 R^1 is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by
20 at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, -SO₂R², C_{0-8} alkyl-C(=O)OH and C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, and where R^2 and R^3 are as described above;

q is an integer from 1-3;

25

R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl, C_{1-6} alkyl-C₃₋₈cycloalkyl, -O- R^2 , -O-C(=O) R^2 , - C_{1-8} alkyl-O- R^{10} , - C_{1-8} alkyl-O-C(=O) R^{10} , - C_{1-8} alkyl-C(=O)OR¹⁰, - C_{1-8} alkyl-O-C(=O)OR¹⁰, - C_{1-8} alkyl-C(=O)NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰R¹⁰, - C_{1-8} alkyl-NR¹⁰C(=O) R^{10} , -SR¹⁰, where R^2 is as set forth above and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached;

10 p is an integer from 0-2;


E is a member selected from the group consisting of a direct link, -O-, -N(- R^{11})-, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group
15 consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups;

Q is a member selected from the group consisting of a direct link, a bivalent
20 C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by
25 from 0 to 5 R^{14} groups;

each R^{14} group is independently a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl-C(=O)OH, C_{0-8} alkyl-C(=O)O- C_{1-8} alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -O- R^2 and -O-C(=O) R^2 , an unsubstituted amino group, a mono- or
30 di-substituted amino group, wherein the substituted amino groups are independently

substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



5 wherein

t is an integer from 0 to 6,

u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


5. A compound of claim 4, wherein:

R^1 and R^8 are each independently a lower alkyl group;

R^{11} is C_3 to C_8 cycloalkyl; and

E and Q are independently a phenylene, a heteroaryl or a heterocyclic group,
5 as set forth above.


6. A compound of formula IIIa:



wherein:


10 p is an integer from 1-2;

E is selected from the group consisting of:




Q is selected from the group consisting of:

and




G is selected from the group consisting of:



5

7. A compound of formula IV:



10 wherein:

R^2 and R^8 is independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the

15 ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group


consisting of N, O and S;

- R¹ and R¹⁴ are independently selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² is as described above;

q is an integer from 0-3;

- R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as set forth above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached;
- and

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein:

5 t is an integer from 0 to 6,

u is the integer 0 or 1, and R¹⁷, R¹⁸, R¹⁹, R²⁰, R²¹, R²², R²³, R²⁴, R²⁵ and R²⁶ are independently selected from the group consisting of H, -OH, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C₁₋₆alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R¹⁸ taken with R¹⁹, R²² taken with either of R²⁴ and R²⁵, and R²⁴ taken with R²⁵, can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

15 with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

8. A compound of claim 7, wherein:

each of R^1 , R^8 and R^{14} is independently selected from the group consisting of hydrogen and C_1 - C_5 alkyl.

5 9. A compound of claim 7, wherein R^8 and R^{14} are each a methyl group, q is 1,

R^1 is a member selected from the group consisting of:


-H , -OH , -NH₂ , -NHCH₂-COOH , -NHSO₂-CH₃ ,

-Br , -COOH , -COOCH₃ , -CF₃ , and -CONH₂

and


R^{11} is a member selected from the group consisting of:

-CH₃ , -CH₂-CH₃ , -CH(-CH₃)₂ , -CH₂COOH




-CH₂CH₂OH , and -C(-CH₃)₃ .

10. A compound of formula V:



5 wherein:

A is a member selected from the group consisting of: R^2 ; $-NR^3R^4$;
 $-C(=O)NR^3R^4$;



10 wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , and R^9 are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4
 15 of such atoms being selected from the group consisting of N, O and S; where R^6 taken with either of R^7 and R^8 , and/or R^7 taken with R^8 , can each form a 5 to 6 membered

heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

- 5 Z is a member selected from the group consisting of a direct link, C₁₋₈alkyl, C₃₋₈cycloalkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₁₋₈carbocyclic aryl, or a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S;

- 10 n is an integer from 0-3;

D is a member selected from the group consisting of a direct link, -CH₂-, -O-, -N(R²)-, -C(=O)-, -S-, -SO₂-, -SO₂-N(R²)-, -N(R²)-SO₂-, -OC(=O)-, -C(=O)O-, -C(=O)-N(R²)- and -N(R²)-C(=O)-, where R² is as described above;

- 15 R¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, halogen, polyhaloalkyl, C₀₋₈alkyl-C(=O)OH, C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, -CN, -NO₂, C₀₋₈alkyl-OH, C₀₋₈alkyl-SH, -C(=O)NR²R³, -O-R² and -O-C(=O)R², an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by
20 at least one member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, polyhaloalkyl, -SO₂R², C₀₋₈alkyl-C(=O)OH and C₀₋₈alkyl-C(=O)O-C₁₋₈alkyl, and where R² and R³ are as described above;


- 25 q is an integer from 0-3;

- R¹¹ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, C₃₋₈cycloalkyl, C₆₋₁₂carbocyclic aryl, C₁₋₆alkylaryl, C₁₋₆alkyl-C₃₋₈cycloalkyl, -O-R², -O-C(=O)R², -C₁₋₈alkyl-O-R¹⁰, -C₁₋₈alkyl-O-C(=O)R¹⁰, -C₁₋₈alkyl-C(=O)OR¹⁰, -C₁₋₈alkyl-O-C(=O)OR¹⁰, -C₁₋₈alkyl-C(=O)NR¹⁰R¹⁰,
30 -C₁₋₈alkyl-NR¹⁰R¹⁰, -C₁₋₈alkyl-NR¹⁰C(=O)R¹⁰, -SR¹⁰, where R² is as described above and R¹⁰ is a member selected from the group consisting of H, C₁₋₈alkyl, C₂₋₈alkenyl, C₂₋₈alkynyl, and wherein when two R¹⁰ groups are present they may be taken together to

form a saturated or unsaturated ring with the atom to which they are both attached;

- each R^{14} group is independently a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$, C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl, -CN, -NO₂, C_{0-8} alkyl-OH, C_{0-8} alkyl-SH, -O- R^2 and -O- $C(=O)R^2$, an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$ and C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,

- u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} ,

can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

- 5 and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


11. A compound of claim 10, wherein each of R¹ and R¹⁴ is independently selected from the group consisting of hydrogen and C₁-C₃alkyl; and R¹¹ is H, or C₃ to C₈ cycloalkyl.

- 10 12. A compound of claim 10, wherein




q is 0;

R¹¹ is -CH₂C(=O)OH;


G is



- 15 A is a member selected from the group consisting of:




Z is a member selected from the group consisting of:



5


n is an integer from 0-2; and

D is a member selected from the group consisting of:



10

13. A compound of formula VI:



wherein:

15 R^6 and R^9 are independently selected from the group consisting of H, $-OH$, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the

ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S;

R^{11} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, C_{1-6} alkylaryl, C_{1-6} alkyl- C_{3-8} cycloalkyl, $-O-R^2$, $-O-C(=O)R^2$, $-C_{1-8}$ alkyl- $O-R^{10}$, $-C_{1-8}$ alkyl- $O-C(=O)R^{10}$, $-C_{1-8}$ alkyl- $C(=O)OR^{10}$, $-C_{1-8}$ alkyl- $O-C(=O)OR^{10}$, $-C_{1-8}$ alkyl- $C(=O)NR^{10}R^{10}$, $-C_{1-8}$ alkyl- $NR^{10}R^{10}$, $-C_{1-8}$ alkyl- $NR^{10}C(=O)R^{10}$, $-SR^{10}$, where R^2 is as described above and R^{10} is a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, and wherein when two R^{10} groups are present they may be taken together to form a saturated or unsaturated ring with the atom to which they are both attached;

X is N or $-CR^{11}$;


p is an integer from 0-3;

E is a member selected from the group consisting of a direct link, $-O-$, $-N(-R^{11})-$, where R^{11} is as defined above, phenylene, a bivalent 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S, wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups;

each R^{14} group is independently a member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, halogen, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$, C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl, $-CN$, $-NO_2$, C_{0-8} alkyl- OH , C_{0-8} alkyl- SH , $-O-R^2$ and $-O-C(=O)R^2$, an unsubstituted amino group, a mono- or di-substituted amino group, wherein the substituted amino groups are independently substituted by at least one member selected from the group consisting of H, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, polyhaloalkyl, C_{0-8} alkyl- $C(=O)OH$ and C_{0-8} alkyl- $C(=O)O-C_{1-8}$ alkyl;

Q is a member selected from the group consisting of a direct link, a bivalent C_{3-8} cycloalkyl group, phenylene, a 5 to 12 member bivalent heteroaryl group containing 1 to 4 heteroatoms selected from the group consisting of N, O and S, and a five to ten membered non-aromatic bivalent heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S wherein said heteroaryl and said non-aromatic heterocyclic ring structure may be independently substituted by from 0 to 5 R^{14} groups;

G is a member selected from the group consisting of: H; -CN; -OR¹⁷;



wherein

t is an integer from 0 to 6,


u is the integer 0 or 1, and R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{22} , R^{23} , R^{24} , R^{25} and R^{26} are independently selected from the group consisting of H, -OH, C_{1-8} alkyl, C_{2-8} alkenyl, C_{2-8} alkynyl, C_{3-8} cycloalkyl, C_{6-12} carbocyclic aryl, a five to ten membered heterocyclic ring system containing 1-4 heteroatoms selected from the group consisting of N, O and S; and C_{1-6} alkylheterocyclic ring system having in the ring system 5 to 10 atoms with 1 to 4 of such atoms being selected from the group consisting of N, O and S; where R^{18} taken with R^{19} , R^{22} taken with either of R^{24} and R^{25} , and R^{24} taken with R^{25} , can each independently form a 5 to 6 membered heterocyclic ring containing from 1 to 4 atoms selected from the group consisting of N, O and S;

with the proviso that when G is H; -CN; -OR¹⁷, either E or Q must contain at least one N atom;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


5

14. A compound of claim 13 having the following structural formula:



15. A compound of claim 14 having the following structural formula:

10



16. A compound of claim 15, wherein:

X is N or CH;

15


R⁹ is a methyl group;

R¹¹ is a member selected from the group consisting of: -C(=O)OCH₂CH₃, -CH(CH₃)₂, -CHCH₃, and -CH₃; and


20

R¹⁴ is a member selected from the group consisting of: -CHCH₃ and -CH₃.

17. A compound selected from the group consisting of:




5



10

and

15



and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

5

18. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of one of claims 1-17.

10

19. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of one of claims 1-17.

15

20. The method of claim 19, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans,

20

thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring
5 the fitting of prosthetic devices.

21. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of one of claims 1-17.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/03225

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 A61K31/4184 C07D401/14 C07D475/04 C07D403/14
C07D235/16 C07D209/10 A61P7/02 A61K31/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 540 051 A (DAIICHI SEIYAKU CO) 5 May 1993 (1993-05-05) the whole document	4-21
Y	WHITLOW, M. ET AL.: "Crystallographic analysis of potent and selective factor Xa inhibitors complexed to bovine trypsin" ACTA CRISTALLOGR., SECT. D: BIOL. CRYSTALLOGR., vol. D55, no. 8, 1999, pages 1395-1404, XP001000519 page 1398	4-21
Y	US 5 849 759 A (ZHAO ZUCHUN ET AL) 15 December 1998 (1998-12-15) the whole document	4-21
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

31 May 2001

Date of mailing of the international search report

13/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/03225

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	IWANOWICZ E J ET AL: "Derivatives of 5-amidine indole as inhibitors of thrombin catalytic activity" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, GB, OXFORD, vol. 6, no. 12, 18 June 1996 (1996-06-18), pages 1339-1344, XP004134837 ISSN: 0960-894X the whole document ----	4-21
A	DE 39 28 177 A (THOMAE GMBH DR K) 28 February 1991 (1991-02-28) page 18, line 45; claim 1 ----	10
A	WO 99 16755 A (HUNGATE RANDALL W ; KOESTER TIMOTHY J (US); BILODEAU MARK T (US); M) 8 April 1999 (1999-04-08) page 31; claim 1 -----	4-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/03225

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0540051 A	05-05-1993	AT 136293 T	15-04-1996
		AU 666137 B	01-02-1996
		AU 2747092 A	06-05-1993
		CA 2081836 A	01-05-1993
		CN 1072677 A,B	02-06-1993
		CN 1168885 A	31-12-1997
		CN 1168886 A,B	31-12-1997
		CZ 284381 B	11-11-1998
		DE 69209615 D	09-05-1996
		DE 69209615 T	09-01-1997
		DK 540051 T	06-05-1996
		ES 2088073 T	01-08-1996
		FI 924932 A	01-05-1993
		GR 3019832 T	31-08-1996
		HK 1002999 A	30-09-1998
		HR 921147 B	30-04-1999
		HR 921147 A	31-10-1995
		HU 65890 A	28-07-1994
		IL 103564 A	06-12-1998
		JP 10291931 A	04-11-1998
		JP 2879718 B	05-04-1999
		JP 5208946 A	20-08-1993
		KR 205152 B	01-07-1999
		MX 9206295 A	01-08-1993
		NO 302948 B	11-05-1998
		NZ 244936 A	26-05-1995
		PL 170312 B	29-11-1996
		RU 2139851 C	20-10-1999
		SK 327692 A	13-04-1999
		US 5962695 A	05-10-1999
		US 5576343 A	19-11-1996
		US 5620991 A	15-04-1997
		US 5866577 A	02-02-1999
		ZA 9208276 A	06-05-1993
US 5849759 A	15-12-1998	AU 700894 B	14-01-1999
		AU 1395697 A	03-07-1997
		CN 1209062 A	24-02-1999
		CZ 9801776 A	16-09-1998
		EP 0865281 A	23-09-1998
		HU 9904066 A	28-03-2000
		WO 9721437 A	19-06-1997
		JP 2000502082 T	22-02-2000
		NO 982606 A	10-08-1998
		PL 327169 A	23-11-1998
		SK 74698 A	02-12-1998
DE 3928177 A	28-02-1991	DE 3911603 A	18-10-1990
		AT 132491 T	15-01-1996
		AU 629324 B	01-10-1992
		AU 5301390 A	11-10-1990
		CA 2014008 A	08-10-1990
		DE 59010017 D	15-02-1996
		DK 392317 T	29-04-1996
		EP 0392317 A	17-10-1990
		ES 2088915 T	01-10-1996
		FI 103044 B	15-04-1999
		GR 3019354 T	30-06-1996

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 01/03225

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3928177 A		HU 53619 A	28-11-1990
		IE 71943 B	12-03-1997
		IL 94049 A	30-05-1994
		JP 1991892 C	22-11-1995
		JP 3063264 A	19-03-1991
		JP 7025739 B	22-03-1995
		KR 171046 B	01-02-1999
		NO 177533 B	26-06-1995
		NZ 233259 A	26-08-1992
		PT 93689 A, B	20-11-1990
		RU 2026861 C	20-01-1995
		US 5541229 A	30-07-1996
		US 5864043 A	26-01-1999
		DD 293581 A	05-09-1991
		ZA 9002695 A	24-12-1991
WO 9916755 A	08-04-1999	AU 9500398 A	23-04-1999
		EP 1017682 A	12-07-2000
		US 6162804 A	19-12-2000